

ECE 4973/5973:

Lecture 12

▼ Image Warping

Samuel Cheng


Slide credit: Noah Snavely, James Thompkin

# Image stitching




Why don't these image line up exactly?

# What is the geometric relationship between these two images?



**Answer: Similarity transformation** (translation, rotation, uniform scale)

What is the geometric relationship between these two images?



What is the geometric relationship between these two images?




**Very important for creating mosaics!**

# Image Warping


- image filtering: change *range* of image

- $g(x) = h(f(x))$



- image warping: change *domain* of image


- $g(x) = f(h(x))$



# Image Warping


- image filtering: change *range* of image

- $g(x) = h(f(x))$



- image warping: change *domain* of image

- $g(x) = f(h(x))$



# Parametric (global) warping

- Examples of parametric warps:



translation



rotation




aspect



# Parametric (global) warping



$\mathbf{p} = (x, y)$



$\mathbf{p}' = (x', y')$

- Transformation  $T$  is a coordinate-changing machine:

$$\mathbf{p}' = T(\mathbf{p})$$

- What does it mean that  $T$  is global?
  - Is the same for any point  $\mathbf{p}$
  - can be described by just a few numbers (parameters)
- Let's consider *linear* xforms (can be represented by a 2D matrix):

$$\mathbf{p}' = \mathbf{T}\mathbf{p} \quad \begin{bmatrix} x' \\ y' \end{bmatrix} = \mathbf{T} \begin{bmatrix} x \\ y \end{bmatrix}$$

# Common linear transformations

- Uniform scaling by  $s$ :




$$\mathbf{S} = \begin{bmatrix} s & 0 \\ 0 & s \end{bmatrix}$$

What is the inverse?

# Common linear transformations

- Rotation by angle  $\theta$  (about the origin)



$$\mathbf{R} = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$$

What is the inverse?

For rotations:

$$\mathbf{R}^{-1} = \mathbf{R}^T$$

# 2x2 Matrices

- What types of transformations can be represented with a 2x2 matrix?

2D mirror about Y axis?

$$\begin{aligned}x' &= -x \\ y' &= y\end{aligned} \quad \mathbf{T} = \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}$$

2D mirror across line  $y = x$ ?

$$\begin{aligned}x' &= y \\ y' &= x\end{aligned} \quad \mathbf{T} = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$$

# 2x2 Matrices

- What types of transformations can be represented with a 2x2 matrix?

2D Translation?

$$x' = x + t_x \quad \text{NO!}$$

$$y' = y + t_y$$

Translation is not a linear operation on 2D coordinates

# All 2D Linear Transformations

- Linear transformations are combinations of ...

- Scale,
- Rotation,
- Shear, and
- Mirror

$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

- Properties of linear transformations:

- Origin maps to origin
- Lines map to lines
- Parallel lines remain parallel
- Ratios are preserved
- Closed under composition


$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} e & f \\ g & h \end{bmatrix} \begin{bmatrix} i & j \\ k & l \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

# Homogeneous coordinates

Trick: add one more coordinate:

$$(x, y) \Rightarrow \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

homogeneous image  
coordinates



Converting *from* homogeneous coordinates

$$\begin{bmatrix} x \\ y \\ w \end{bmatrix} \Rightarrow (x/w, y/w)$$

# Translation

- Solution: homogeneous coordinates to the rescue


$$\mathbf{T} = \begin{bmatrix} 1 & 0 & t_x \\ 0 & 1 & t_y \\ 0 & 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & t_x \\ 0 & 1 & t_y \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} = \begin{bmatrix} x + t_x \\ y + t_y \\ 1 \end{bmatrix}$$



# Affine transformations

$$\mathbf{T} = \begin{bmatrix} 1 & 0 & t_x \\ 0 & 1 & t_y \\ 0 & 0 & 1 \end{bmatrix}$$



any transformation with  
last row  $[0 \ 0 \ 1]$  we call an  
*affine* transformation

$$\begin{bmatrix} a & b & c \\ d & e & f \\ 0 & 0 & 1 \end{bmatrix}$$

# Basic affine transformations

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & t_x \\ 0 & 1 & t_y \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

Translate

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} s_x & 0 & 0 \\ 0 & s_y & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

Scale

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} \cos \theta & -\sin \theta & 0 \\ \sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

2D *in-plane* rotation

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} 1 & sh_x & 0 \\ sh_y & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

Shear

# Affine Transformations

- Affine transformations are combinations of ...

- Linear transformations, and
- Translations

$$\begin{bmatrix} x' \\ y' \\ w \end{bmatrix} = \begin{bmatrix} a & b & c \\ d & e & f \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ w \end{bmatrix}$$


- Properties of affine transformations:
  - Origin does not necessarily map to origin
  - Lines map to lines
  - Parallel lines remain parallel
  - Ratios are preserved
  - Closed under composition

# Is this an affine transformation?



# Where do we go from here?

$$\begin{bmatrix} a & b & c \\ d & e & f \\ 0 & 0 & 1 \end{bmatrix}$$



what happens when we  
mess with this row?

affine transformation


# Projective Transformations aka Homographies aka Planar Perspective Maps

$$\mathbf{H} = \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & 1 \end{bmatrix}$$

Called a *homography*  
(or *planar perspective map*)



# Image warping with homographies



# Homographies





# Projective Transformations

- Projective transformations ...
    - Affine transformations, and
    - Projective warps
- $$\begin{bmatrix} x' \\ y' \\ w' \end{bmatrix} = \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} \begin{bmatrix} x \\ y \\ w \end{bmatrix}$$
- Properties of projective transformations:
    - Origin does not necessarily map to origin
    - Lines map to lines
    - Parallel lines do not necessarily remain parallel
    - Ratios are not preserved
    - Closed under composition

# Transforming coordinate vs transforming bases


- In earlier examples, we have fixed bases but transforming coordinate. Note that

$$- [b_1 b_2] \left( T \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} \right) = ([b_1 b_2] T) \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = ([b_1 T, b_2 T]) \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

– It is equivalent to

# Summary

- Image warping vs filtering
  - $I(f(x))$  vs  $f(I(x))$
- Global warping
- Linear vs affine
  - $Ax$  vs  $Ax + b$
- Homogeneous coordinate vs Cartesian coordinate
- Affine vs projective transform



Transform	DOF	Preserve
Translation	2	Orientation + p(rigid)
Rotation	1	Center of Mass + p(rigid)
Rigid (Euclidean)	3	Length + p(similarity)
Similarity	4	Angle + p(affine)
Affine	6	Parallelism + p(projective)
Projective	8	Straight line